



Planning My Routine

Use this Treatment Calendar to help you stay organized.



“Treatment means there are possibilities.”

– Dinah, former patient who received EMPLICITI

Use this resource to help you understand and plan your EMPLICITI® (elotuzumab) dosing schedule for:

EMPLICITI + POMALYST® (pomalidomide) + dexamethasone (EPd)

EMPLICITI + REVLIMID® (lenalidomide) + dexamethasone (ERd)

What is EMPLICITI?

EMPLICITI is a prescription medicine used to treat multiple myeloma in combination with the medicines:

- REVLIMID and dexamethasone in adults who have received one to three prior treatments for their multiple myeloma.
- POMALYST and dexamethasone in adults who have received at least two prior treatments including REVLIMID and a proteasome inhibitor.

It is not known if EMPLICITI, REVLIMID, or POMALYST is safe and effective in children.

Important Safety Information for EMPLICITI®, REVLIMID®, and POMALYST®

EMPLICITI in combination with REVLIMID and dexamethasone or in combination with POMALYST and dexamethasone may cause the following serious side effects: Infusion reactions, infections, risk of developing new cancers (malignancies), and liver problems. **There are also other serious risks (Boxed WARNINGS) associated with REVLIMID and POMALYST. REVLIMID and POMALYST can cause harm to unborn babies and blood clots. REVLIMID can also cause low blood counts. REVLIMID and POMALYST are only available through the Lenalidomide REMS or POMALYST REMS® programs.** More information on the REMS programs is available at www.CelgeneRiskManagement.com or by calling 1-888-423-5436. **For more safety information, please read the Important Safety Information throughout and on pages 12-20.** Your healthcare team will work with you to manage any side effects you may experience throughout your treatment with EMPLICITI with REVLIMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone.

Please read Patient Information in full Prescribing Information for EMPLICITI. Please read Medication Guide in full Prescribing Information for REVLIMID and POMALYST, including Boxed WARNINGS.

Emplaciti.
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

How will I receive EMPLICITI® (elotuzumab)?

EMPLICITI is given through an intravenous (IV) infusion by your healthcare team. An IV infusion is when medicine is given directly into the bloodstream through a vein—usually in the arm, hand, or through an IV port.



If you miss your EMPLICITI infusion, call your healthcare team right away to schedule another time to get an infusion.

EMPLICITI keeps working between your treatments to help your immune system fight the disease. That's why it is very important to keep every appointment on the day it is scheduled. To make the most of your treatment plan, you should continue receiving EMPLICITI for as long as your healthcare provider prescribes the medication.

Important Safety Information (continued)

Before you receive EMPLICITI, tell your healthcare provider about all of your medical conditions, including if you:

- have an infection
- are pregnant or plan to become pregnant. It is not known if EMPLICITI may harm your unborn baby. However, REVLIMID® (lenalidomide) and POMALYST® (pomalidomide) may cause birth defects or death of an unborn baby.
- **Before receiving EMPLICITI with REVLIMID and dexamethasone, or EMPLICITI with POMALYST and dexamethasone, females and males must agree to the instructions in the Lenalidomide REMS program or the POMALYST REMS® program, depending on which combination your doctor has prescribed for you. The programs have specific requirements about birth control (contraception), pregnancy testing, blood donation, and sperm donation that you need to know. Talk to your healthcare provider to learn more about REVLIMID or POMALYST.**

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.

Emplificiti.
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

A treatment plan that can fit into your routine

Your treatment schedule will depend on your treatment plan and the other medications you will be taking in combination with EMPLICITI® (elotuzumab).



Your EMPLICITI treatment schedule is divided into cycles that are 28 days (4 weeks) long.

Within each treatment cycle is a rest period. This is the number of days that you will not be receiving any EMPLICITI with REVLIMID® (lenalidomide) and dexamethasone, or with POMALYST® (pomalidomide) and dexamethasone. Rest periods are usually at the end of each treatment cycle.

Important Safety Information (continued)

Before you receive EMPLICITI, tell your healthcare provider about all of your medical conditions, including if you: (continued)

- are breastfeeding or plan to breastfeed. It is not known if EMPLICITI passes into breast milk. You should not breastfeed during treatment with EMPLICITI with REVLIMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone.
- **Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Serious side effects that can occur with EMPLICITI treatment are:

Infusion reactions

- Infusion reactions can happen during your infusion or within 24 hours after your infusion of EMPLICITI. Your healthcare provider will give you medicines before each infusion of EMPLICITI to help reduce the risk of an infusion reaction.
- If you have an infusion reaction while receiving EMPLICITI, your healthcare provider will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction your healthcare provider may stop your treatment completely.
- Tell your healthcare provider or get medical help right away if you have any of these symptoms after your infusion with EMPLICITI: fever, chills, rash, chest pain, trouble breathing, dizziness, light-headedness.

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.

Empliciti.
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

Taking your oral dexamethasone dose

On days that you **are** receiving EMPLICITI® (elotuzumab), take your dexamethasone dose:

- by mouth
- exactly as prescribed
- between 3 and 24 hours before your EMPLICITI infusion treatment

On days that you **are not** receiving EMPLICITI, take your dexamethasone dose:

- by mouth
- exactly as prescribed
- according to your treatment schedule

Keep in mind that the dose of oral dexamethasone may vary based on whether EMPLICITI is given that day. If you are receiving EMPLICITI with POMALYST® (pomalidomide) + dexamethasone, your dose of dexamethasone may also vary based on your age. Be sure to follow the instructions of your healthcare provider.

Important Safety Information (continued)

Serious side effects that can occur with EMPLICITI treatment are: (continued)

Infections

- Those receiving EMPLICITI with REVLIMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone may develop infections; some can be serious.
- Tell your healthcare provider right away if you have any of the signs and symptoms of an infection, including: fever, flu-like symptoms, cough, shortness of breath, burning with urination, or a painful skin rash.

Risk of new cancers (malignancies)

- Those receiving EMPLICITI with REVLIMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone have a risk of developing new cancers.
- Talk with your healthcare provider about your risk of developing new cancers if you receive EMPLICITI.
- Your healthcare provider will check you for new cancers during your treatment with EMPLICITI.

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.

Empliciti.
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

EPd

EMPLICITI + POMALYST® (pomalidomide) + dexamethasone

Your EMPLICITI treatment schedule is divided into cycles that are 28 days (4 weeks) long. A cycle is a set number of days you are on treatment and also includes the time you spend resting between treatments.

EMPLICITI is usually given 1 time every week for cycles 1 and 2 (28-day cycles). After the first 2 cycles, **EMPLICITI is given once every 4 weeks.**

Before each infusion, your healthcare provider will give you medicines to help reduce the risk of an infusion reaction.

Take your dose of dexamethasone exactly as prescribed. Keep in mind the dose of oral dexamethasone may vary based on your age and whether EMPLICITI is given that day.

CYCLES 1 & 2 (28 DAYS EACH)						CYCLES 3 & UP (28 DAYS EACH)					
EMPLICITI	DAY 1		DAY 8		DAY 15		DAY 22				
POMALYST	DAYS 1-21										
oral dexamethasone	DAY 1		DAY 8		DAY 15		DAY 22				
											DAYS 23-28 REST PERIOD

EMPLICITI	DAY 1										
POMALYST	DAYS 1-21										
oral dexamethasone	DAY 1		DAY 8		DAY 15		DAY 22				
											DAYS 23-28 REST PERIOD

How long does each EMPLICITI infusion take?

The length of each EMPLICITI infusion will depend on your body weight and how many times you have received EMPLICITI in the past. If infusion reactions occur or become worse, it may take more time to receive your EMPLICITI infusion.

Important Safety Information (continued)

Serious side effects that can occur with EMPLICITI treatment are: (continued)

Liver problems

- EMPLICITI may cause liver problems. Your healthcare provider will do blood tests to check your liver during treatment with EMPLICITI.
- Tell your healthcare provider if you have signs and symptoms of liver problems, including: tiredness, weakness, loss of appetite, yellowing of your skin or eyes, color changes in your stools, confusion, or swelling of the stomach area.

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.



ERd

EMPLICITI + REVLIMID® (lenalidomide) + dexamethasone

Your EMPLICITI treatment schedule is divided into cycles that are 28 days (4 weeks) long. A cycle is a set number of days you are on treatment and also includes the time you spend resting between treatments.

EMPLICITI is usually given 1 time every week for cycles 1 and 2 (28-day cycles), and **1 time every 2 weeks** for cycles 3 and up (28-day cycles) when used with REVLIMID and dexamethasone.

Before each infusion, your healthcare provider will give you medicines to help reduce the risk of an infusion reaction.

Take your dose of dexamethasone exactly as prescribed. Keep in mind the dose of oral dexamethasone may vary based on whether EMPLICITI is given that day.

CYCLES 1 & 2 (28 DAYS EACH)							CYCLES 3 & UP (28 DAYS EACH)									
EMPLICITI	DAY 1		DAY 8		DAY 15		DAY 22			DAY 1			DAY 15			DAYS 23-28 REST PERIOD
REVLIMID	DAYS 1-21									DAYS 1-21						
oral dexamethasone	DAY 1		DAY 8		DAY 15		DAY 22			DAY 1		DAY 8		DAY 15	DAY 22	

How long does each EMPLICITI infusion take?

The length of each EMPLICITI infusion will depend on your body weight and how many times you have received EMPLICITI in the past. If infusion reactions occur or become worse, it may take more time to receive your EMPLICITI infusion.

Important Safety Information (continued)

The most common side effects of EMPLICITI when used with REVLIMID and dexamethasone include:

- fatigue
- diarrhea
- fever
- constipation
- cough
- numbness, weakness, tingling, or burning pain in your arms or legs
- sore throat or runny nose
- upper respiratory tract infection
- decreased appetite
- pneumonia

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.



Create a Treatment Calendar

We've provided you with blank calendars to help you organize your EMPLICITI® (elotuzumab) dosing schedule and plan your routine. Your treatment schedule will depend on your treatment plan and the other medications you will be taking in combination with EMPLICITI.

Be sure to work with your doctor to fill out your exact schedule and dosage. Together, you can add the following information:

Dates & Days

Fill in the dates and day numbers (1-28) of your treatment cycle with EMPLICITI.

Treatments

Check the boxes of the treatments you'll be receiving each day. Feel free to write in other information, like your doses and appointment times.

Rest Periods

Include the number of days within each cycle that you will not be receiving any EMPLICITI (**E**) with REVLIMID (**R**) and dexamethasone (**D**), or with POMALYST (**P**) and dexamethasone (**D**).

Month _____

Don't forget to take your dexamethasone dose by mouth, 3-24 hours before your EMPLICITI treatments!

E = EMPLICITI® (elotuzumab) R = REVLIMID® (lenalidomide) P = POMALYST® (pomalidomide) D = dexamethasone

<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D
date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____
<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D
date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____
<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D
date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____
<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or P <input type="checkbox"/> D
date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____	date _____ day _____

Important Safety Information (continued)

The most common side effects of EMPLICITI when used with POMALYST and dexamethasone include:

- constipation
- high blood sugar

These are not all of the possible side effects of EMPLICITI. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for [EMPLICITI](#). Please read Medication Guide in full Prescribing Information for [REVLIMID](#) and [POMALYST](#), including Boxed WARNINGS.

Emplciti.
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

Your Treatment Calendar

Month _____

Don't forget to take your dexamethasone dose by mouth, 3-24 hours before your EMPLICITI treatments!

E = EMPLICITI® (elotuzumab) **R** = REVLIMID® (lenalidomide) **P** = POMALYST® (pomalidomide) **D** = dexamethasone

<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for **EMPLICITI**. Please read Medication Guide in full Prescribing Information for **REVLIMID** and **POMALYST**, including Boxed WARNINGS.

EmPLICITI
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

Your Treatment Calendar

Month _____

Don't forget to take your dexamethasone dose by mouth, 3-24 hours before your EMPLICITI treatments!

E = EMPLICITI® (elotuzumab) **R** = REVLIMID® (lenalidomide) **P** = POMALYST® (pomalidomide) **D** = dexamethasone

<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for **EMPLICITI**. Please read Medication Guide in full Prescribing Information for **REVLIMID** and **POMALYST**, including Boxed WARNINGS.

EmPLICITI
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

Your Treatment Calendar

Month _____

Don't forget to take your dexamethasone dose by mouth, 3-24 hours before your EMPLICITI treatments!

E = EMPLICITI® (elotuzumab) **R** = REVLIMID® (lenalidomide) **P** = POMALYST® (pomalidomide) **D** = dexamethasone

<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day
<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D	<input type="checkbox"/> E <input type="checkbox"/> R or <input type="checkbox"/> P <input type="checkbox"/> D
_____ day	_____ day	_____ day	_____ day	_____ day	_____ day	_____ day

Please see additional Important Safety Information throughout and on pages 12-20, and read Patient Information in full Prescribing Information for **EMPLICITI**. Please read Medication Guide in full Prescribing Information for **REVLIMID** and **POMALYST**, including Boxed WARNINGS.

EmPLICITI
(elotuzumab)
FOR INJECTION
FOR INTRAVENOUS USE
300 MG & 400 MG VIALS

What is EMPLICITI® (elotuzumab)?

EMPLICITI is a prescription medicine used to treat multiple myeloma in combination with the medicines:

- REVLMID® (lenalidomide) and dexamethasone in adults who have received one to three prior treatments for their multiple myeloma.
- POMALYST® (pomalidomide) and dexamethasone in adults who have received at least two prior treatments including REVLMID and a proteasome inhibitor.

What is REVLMID?

REVLMID is a prescription medicine used to treat adults with multiple myeloma in combination with the medicine dexamethasone, or as maintenance treatment after autologous hematopoietic stem cell transplantation (a type of stem cell transplant that uses your own stem cells). REVLMID should not be used to treat people who have chronic lymphocytic leukemia (CLL) unless they are participants in a controlled clinical trial.

What is POMALYST?

POMALYST is a prescription medicine, taken along with the medicine dexamethasone, used to treat adults with multiple myeloma who have previously received at least 2 medicines to treat multiple myeloma, including a proteasome inhibitor and lenalidomide, and whose disease has become worse during treatment or within 60 days of finishing the last treatment.

It is not known if EMPLICITI, REVLMID, or POMALYST is safe and effective in children.

Important Safety Information

WARNINGS FOR REVLMID: Risk to unborn babies, risk of low blood counts and blood clots.

WARNINGS FOR POMALYST: Risk to unborn babies, and blood clots.

What is the most important information I should know about REVLMID & POMALYST?

Before you begin taking REVLMID or POMALYST, you must read and agree to all of the instructions in the Lenalidomide REMS or POMALYST REMS® program. Before prescribing REVLMID or POMALYST, your healthcare provider (HCP) will explain the Lenalidomide REMS or POMALYST REMS program to you and have you sign the Patient-Physician Agreement Form.




Important Safety Information (continued)

What is the most important information I should know about REVLMID® (lenalidomide) & POMALYST® (pomalidomide)? (continued)

REVLMID & POMALYST can cause serious side effects, including:

- **Possible birth defects (deformed babies) or death of an unborn baby.** Females who are pregnant or plan to become pregnant must not take REVLMID or POMALYST.
 - **REVLMID & POMALYST are similar to the medicine thalidomide (THALOMID®)**, which is known to cause severe life-threatening birth defects. REVLMID & POMALYST have not been tested in pregnant females. REVLMID & POMALYST have harmed unborn animals in animal testing.
- **Females must not get pregnant:**
 - o For at least 4 weeks before starting REVLMID or POMALYST
 - o While taking REVLMID or POMALYST
 - o During any breaks (interruptions) in your treatment with REVLMID or POMALYST
 - o For at least 4 weeks after stopping REVLMID or POMALYST
- **Females who can become pregnant:**
 - o Will have pregnancy tests weekly for 4 weeks, then every 4 weeks if your menstrual cycle is regular, or every 2 weeks if your menstrual cycle is irregular. If you miss your period or have unusual bleeding, you will need to have a pregnancy test and receive counseling.
 - o Must agree to use 2 acceptable forms of effective birth control at the same time, for at least 4 weeks before, while taking, during any breaks (interruptions) in treatment, and for at least 4 weeks after stopping REVLMID or POMALYST.
 - o Talk with your healthcare provider to find out about options for acceptable forms of birth control that you may use to prevent pregnancy during and after treatment with REVLMID or POMALYST.
- **If you become pregnant while taking REVLMID or POMALYST, stop taking it right away and call your healthcare provider.** If your healthcare provider is not available, you can call Celgene Customer Care Center at 1-888-423-5436. Healthcare providers and patients should report all cases of pregnancy to:
 - o FDA MedWatch at 1-800-FDA-1088
 - o Celgene Corporation at 1-888-423-5436



Important Safety Information (continued)

What is the most important information I should know about REVLMID® (lenalidomide) & POMALYST® (pomalidomide)? (continued)

There is a pregnancy exposure registry that monitors the outcomes of females who take REVLMID or POMALYST during pregnancy, or if their male partner takes REVLMID or POMALYST and they are exposed during pregnancy. You can enroll in this registry by calling Celgene Corporation at the phone number listed on page 13.

– **REVLMID & POMALYST can pass into human semen:**

- o Males, including those who have had a vasectomy, must always use a latex or synthetic condom during any sexual contact with a pregnant female or a female that can become pregnant while taking REVLMID or POMALYST, during any breaks (interruptions) in your treatment with REVLMID or POMALYST, and for 4 weeks after stopping REVLMID or POMALYST.
- o Do not have unprotected sexual contact with a female who is or could become pregnant. Tell your healthcare provider if you do have unprotected sexual contact with a female who is or could become pregnant.
- o Do not donate sperm while taking REVLMID or POMALYST, during any breaks (interruptions) in your treatment, and for 4 weeks after stopping REVLMID or POMALYST. If a female becomes pregnant with your sperm, the baby may be exposed to REVLMID or POMALYST and may be born with birth defects.

Men, if your female partner becomes pregnant, you should call your healthcare provider right away.

- **Do not donate blood** while you take REVLMID or POMALYST, during any breaks (interruptions) in your treatment, and for 4 weeks after stopping REVLMID or POMALYST. If someone who is pregnant gets your donated blood, her baby may be exposed to REVLMID or POMALYST and may be born with birth defects.

- **Low white blood cells (neutropenia) and low platelets (thrombocytopenia).** REVLMID causes low white blood cells and low platelets in most people. You may need a blood transfusion or certain medicines if your blood counts drop too low. Your healthcare provider should check your blood counts often, especially during the first several months of treatment with REVLMID, and then at least monthly. Tell your healthcare provider if you develop any bleeding or bruising during treatment with REVLMID.



Important Safety Information (continued)

What is the most important information I should know about REVLMID® (lenalidomide) & POMALYST® (pomalidomide)? (continued)

- **Blood clots in your arteries, veins, and lungs, heart attack, and stroke can happen if you take REVLMID or POMALYST.**

- Most people who take REVLMID or POMALYST will also take a blood thinner medicine to help prevent blood clots.
- Before taking REVLMID or POMALYST, tell your healthcare provider:
 - o If you have had a blood clot in the past.
 - o If you have high blood pressure, smoke, or if you have been told you have a high level of fat in your blood (hyperlipidemia).
 - o About all the medicines you take. Certain other medicines can also increase your risk for blood clots.

Call your healthcare provider or get medical help right away if you get any of the following during treatment with REVLMID or POMALYST:

- smoke cigarettes (POMALYST may not work as well in people who smoke)
- have liver problems
- have kidney problems or receive hemodialysis treatment
- have thyroid problems
- have an infection
- have had a serious skin rash with thalidomide treatment. You should not take REVLMID.
- are lactose intolerant. REVLMID contains lactose.
- Are pregnant or plan to become pregnant. It is not known if EMPLICITI may harm your unborn baby. However, REVLMID & POMALYST may cause birth defects or death of an unborn baby.
- are breastfeeding. Do not breastfeed during treatment with EMPLICITI and REVLMID and dexamethasone or EMPLICITI and POMALYST and dexamethasone.
- **Tell your HCP about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REVLMID and other medicines or POMALYST and other medicines may affect each other, causing serious side effects. Talk with your HCP before taking any new medicines.
- Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist.

Get emergency medical help right away if you develop any of the following signs or symptoms during treatment with REVLMID or POMALYST:

- o swelling of your lips, mouth, tongue, or throat
- o a very fast heartbeat
- o trouble breathing or swallowing
- o you feel dizzy or faint
- o raised red areas on your skin (hives)



Important Safety Information (continued)

Who should not take REVLMID® (lenalidomide) or POMALYST® (pomalidomide)?

Do not take REVLMID or POMALYST if you:

- Are pregnant, plan to become pregnant, or become pregnant during treatment with REVLMID or POMALYST. **See “What is the most important information I should know about REVLMID & POMALYST?”**
- Are allergic to lenalidomide or pomalidomide or any of the ingredients in REVLMID or POMALYST.


What should I tell my healthcare provider (HCP) before taking EMPLICITI, REVLMID, or POMALYST?

Before you take EMPLICITI, REVLMID, or POMALYST, tell your healthcare provider about all of your medical conditions, including if you:

- smoke cigarettes (POMALYST may not work as well in people who smoke)
- have liver problems
- have kidney problems or receive hemodialysis treatment
- have thyroid problems
- have an infection
- have had a serious skin rash with thalidomide treatment. You should not take REVLMID.
- are lactose intolerant. REVLMID contains lactose.
- Are pregnant or plan to become pregnant. It is not known if EMPLICITI may harm your unborn baby. However, REVLMID & POMALYST may cause birth defects or death of an unborn baby.
- are breastfeeding. Do not breastfeed during treatment with EMPLICITI and REVLMID and dexamethasone or EMPLICITI and POMALYST and dexamethasone.
- **Tell your HCP about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. REVLMID and other medicines or POMALYST and other medicines may affect each other, causing serious side effects. Talk with your HCP before taking any new medicines.
- Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist.

What are the possible side effects of EMPLICITI, REVLMID, and POMALYST?

- **See “What is the most important information I should know about REVLMID & POMALYST?”**
- **EMPLICITI, REVLMID, and POMALYST can cause serious side effects, including:**
 - **Increased risk of death in people who have chronic lymphocytic leukemia (CLL).** People with CLL who take REVLMID have an increased risk of death compared with people who take the medicine chlorambucil. REVLMID may cause you to have serious heart problems that can lead to death, including atrial fibrillation, heart attack, or heart failure. You should not take REVLMID if you have CLL unless you are participating in a controlled clinical trial.



Important Safety Information (continued)

How should I take REVLMID® (lenalidomide) or POMALYST® (pomalidomide)?

Take REVLMID or POMALYST exactly as prescribed and follow all the instructions of the Lenalidomide REMS & POMALYST REMS® program.

- Swallow REVLMID or POMALYST capsules whole with water 1 time a day.


Do not break, chew, or open your capsules.

- REVLMID or POMALYST may be taken with or without food.
- Take REVLMID or POMALYST at the same time each day.
- If you are on hemodialysis, take POMALYST after hemodialysis, on hemodialysis days.
- Do not open REVLMID or POMALYST capsules or handle them any more than needed. If you touch a broken REVLMID or POMALYST capsule or the medicine in the capsule, wash the area of your body right away with soap and water.

- If you miss a dose of REVLMID or POMALYST and it has been less than 12 hours since your regular time, take REVLMID or POMALYST as soon as you remember. If it has been more than 12 hours, just skip your missed dose. **Do not** take 2 doses at the same time.
- If you take too much REVLMID or POMALYST, call your HCP right away.
- **Do not share REVLMID & POMALYST with other people.** It may cause birth defects and other serious problems.

What are the possible side effects of EMPLICITI, REVLMID, and POMALYST?


- **See “What is the most important information I should know about REVLMID & POMALYST?”**
- **EMPLICITI, REVLMID, and POMALYST can cause serious side effects, including:**
 - **Increased risk of death in people who have chronic lymphocytic leukemia (CLL).** People with CLL who take REVLMID have an increased risk of death compared with people who take the medicine chlorambucil. REVLMID may cause you to have serious heart problems that can lead to death, including atrial fibrillation, heart attack, or heart failure. You should not take REVLMID if you have CLL unless you are participating in a controlled clinical trial.



Important Safety Information (continued)

What are the possible side effects of EMPLICITI, REVLMID® (lenalidomide), and POMALYST® (pomalidomide)? (continued)


- **Low white blood cells (neutropenia), low platelets (thrombocytopenia), and low red blood cells (anemia) are common with POMALYST, but can also be serious.** You may need a blood transfusion or certain medicines if your blood counts drop too low. Your blood counts should be checked by your healthcare provider (HCP) weekly for the first 8 weeks of treatment and monthly after that.
- **Severe liver problems, including liver failure and death.** Your HCP should do blood tests to check your liver function during your treatment with EMPLICITI, REVLMID, and POMALYST. Tell your HCP right away if you develop any of the following symptoms of liver problems: yellowing of your skin or the white parts of your eyes (jaundice); dark or brown (tea-colored) urine; color changes in your stool; pain or swelling on the upper right side of your stomach area (abdomen); confusion; bleeding or bruising more easily than normal, or feeling very tired.
- **Infusion Reactions.** Infusion reactions can happen during your infusion or within 24 hours after your infusion of EMPLICITI. Your healthcare provider will give you medicines before each infusion of EMPLICITI to help reduce the risk of an infusion reaction. If you have an infusion reaction while receiving EMPLICITI, your healthcare provider will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction your healthcare provider may stop your treatment completely. Tell your healthcare provider or get medical help right away if you have any of these symptoms after your infusion with EMPLICITI: fever, chills, rash, chest pain, trouble breathing, dizziness, or light-headedness.
- **Infections.** Those receiving EMPLICITI with REVLMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone may develop infections; some can be serious. Tell your healthcare provider right away if you have any of the signs and symptoms of an infection, including: fever, flu-like symptoms, cough, shortness of breath, burning with urination, or a painful skin rash. **Severe allergic and severe skin reactions** can happen with REVLMID & POMALYST and may cause death.



Important Safety Information (continued)

What are the possible side effects of EMPLICITI, REVLMID® (lenalidomide), and POMALYST® (pomalidomide)? (continued)

- **Dizziness and confusion.** Avoid taking other medicines that may cause dizziness and confusion during treatment with POMALYST. Avoid situations that require you to be alert until you know how POMALYST affects you.
- **Nerve damage.** Stop taking POMALYST and call your HCP if you develop numbness, tingling, pain, or a burning sensation in your hands, legs, or feet.
- **Risk of new cancers (malignancies).** New cancers, including certain blood cancers (acute myelogenous leukemia or AML) have been seen in people who received POMALYST. Those receiving EMPLICITI with REVLMID and dexamethasone or EMPLICITI with POMALYST and dexamethasone have a risk of developing new cancers. Your healthcare provider will check you for new cancers during your treatment with EMPLICITI with REVLMID and dexamethasone, or EMPLICITI with POMALYST and dexamethasone. Talk with your HCP about your risk of developing new cancers.
- **Tumor Lysis Syndrome (TLS).** TLS is caused by the fast breakdown of cancer cells. TLS can cause kidney failure and the need for dialysis treatment, abnormal heart rhythm, seizure, and sometimes death. Your HCP may do blood tests to check you for TLS.
- **Worsening of your tumor (tumor flare reaction).** Tell your healthcare provider if you get any of these symptoms of tumor flare reaction while taking REVLMID: tender, swollen lymph nodes; low-grade fever, pain, or rash.
- **Thyroid problems.** Your healthcare provider may check your thyroid function before you start taking REVLMID and during treatment with REVLMID.
- **Risk of early death in MCL.** In people who have Mantle Cell Lymphoma (MCL), there may be a risk of dying sooner (early death) when taking REVLMID. Talk with your healthcare provider about any concerns and possible risk factors.
- The most common side effects of EMPLICITI when used with REVLMID and dexamethasone include fatigue, diarrhea, fever, constipation, cough, numbness, weakness, tingling, or burning pain in your arms or legs, sore throat or runny nose, upper respiratory tract infection, decreased appetite, and pneumonia.
- The most common side effects of EMPLICITI when used with POMALYST and dexamethasone include constipation and high blood sugar.



Important Safety Information (continued)

What are the possible side effects of EMPLICITI, REVLMID® (lenalidomide), and POMALYST® (pomalidomide)? (continued)

- The most common side effects of REVLMID include diarrhea, rash, nausea, constipation, tiredness or weakness, fever, itching, swelling of your arms, hands, legs, feet and skin, sleep problems (insomnia), headache, muscle cramps or spasms, shortness of breath, cough, sore throat, and other symptoms of a cold, upper respiratory tract infection or bronchitis, inflammation of the stomach and intestine (“stomach flu”), nose bleed, shaking or trembling (tremor), joint aches, and pain in your back or stomach area (abdomen).
- The most common side effects of POMALYST include tiredness and weakness, constipation, nausea, diarrhea, shortness of breath, upper respiratory tract infection, back pain, and fever.
- These are not all the possible side effects of EMPLICITI, REVLMID, and POMALYST. Your HCP may tell you to decrease your dose, temporarily stop or permanently stop taking REVLMID or POMALYST if you develop certain serious side effects during treatment. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.





Planning My Routine

Find more information about starting your
EMPLICITI® (elotuzumab) treatment routine and how
to get more blank calendars at EMPLICITI.com.



EMPLICITI® and its associated logo is a registered trademark of Bristol-Myers Squibb Company.
POMALYST®, POMALYST REMS®, REVLIMID®, and THALOMID® are registered trademarks of
Celgene Corporation, a Bristol Myers Squibb company.